

Original Communication

Aggression and violence towards health care providers – A problem in Turkey? ☆

Hayri Erkol MD (Professor) ^a, Mira R. Gökdoğan PhD ^{b,*},
Zerrin Erkol MD (Associate Professor) ^c, Bora Boz MD (Assistant Professor) ^d

^a Abant İzzet Baysal University, İzzet Baysal Faculty of Medicine, Department of General Surgery, Bolu, Turkey

^b Istanbul University, Institute of Forensic Sciences, Department of Medical Science, Istanbul, Turkey

^c Abant İzzet Baysal University, İzzet Baysal Faculty of Medicine, Department of Legal Medicine, Bolu, Turkey

^d Pamukkale University, Faculty of Medicine, Department of Legal Medicine, Denizli, Turkey

Received 28 November 2006; received in revised form 19 February 2007; accepted 5 March 2007

Available online 12 July 2007

Abstract

Health care providers are increasingly concerned about the escalating incidence of verbal and physical abuse to healthcare staff. Factors, such as long wait in hospital areas, which lead to client frustration over an inability to obtain needed services promptly, are influencing these situations. Nonetheless, incidents of this nature can cause immense psychological harm as well as physical damage among medical employees.

The current study aimed to ascertain from staff members aggressive experiences in the workplace, and the effects on the individual. The results of this study mirrored those of similar surveys in Turkey. Non-reporting was revealed as a major problem. Respondents believed they were treated less seriously than similar incidents involving private citizens. Accordingly, staff criticized hospital managers, the police, and the courts for their attitude about assaults towards hospital employees. They reported feeling vulnerable to abuse and there was a general desire for training in preventing and protection. These include teaching staff breakaway techniques, increasing the number of trained security officers on duty, issuing personal alarms, and encouraging staff to officially report all incidents.

© 2007 Elsevier Ltd and FFLM. All rights reserved.

Keywords: Aggression; Violence; Health care providers; Workplace; Emergency department

1. Introduction

Health reports provide an overview of the health of a country's population and the main factors related to it. Based on international comparisons, they present a summary assessment of what has been achieved so far and what could be improved in the future. A special case of comparison is when each hospital is given a rank order. Although useful as summary measures, ranks can be misleading and should be interpreted with caution. Also, when used to give

an assessment of trends, ranks can hide quite important changes within an individual hospital.^{1–5}

Based on information through mediums such as the Internet, the expectation of patients and their relatives is therefore extremely high, or on the contrary apprehensive, when expecting, wanting, or requiring treatment.⁶ This attitude can easily change into anger, rage, or violence facing realities of clinical practice in emergency departments.⁷ Factors, such as long wait in hospital environments, which lead to client frustration over an inability to obtain needed services promptly, are influencing these situations.⁸ Nonetheless, incidents of this nature can inflict immense psychological harm as well as physical damage among medical employees.⁹ How serious or otherwise this is treated by the victim very much depends on the circumstances and

☆ The results of this article were presented at the IV. Regional Trauma and Emergency Congress, 25–28 September 2003, Kayseri, Turkey.

* Corresponding author. Mobile: +90 532 739 94 02; fax: +90 212 588 00 11.

E-mail address: miragok@istanbul.edu.tr (M.R. Gökdoğan).

the assailant.¹⁰ The current study aimed to ascertain from health care employees aggressive and violent experiences in the workplace, and the effects on the individual. Recommendations are made to increase the awareness of aggressive behavior by setting policies and appropriate, regularly updated training for all health care providers.

2. Materials and methods

For the purpose of this study, a 15-item questionnaire was designed and confidentially offered to physicians, nurses/midwives, and attendants, in emergency departments. Attendants are patient care assistants with handling and lifting duties. The detailed survey regarding aggressive and violent behavior was conducted at the University Hospital, the State Hospital, the Hospital of Gynecology and Pediatrics, and at the Health Clinic, in the province of Bolu, Turkey, over a 5-year period (1998–2003). The hospital's catchments area is largely urban working class and includes many communities, according to their different socio-economic levels. The aims of the survey included determining the following factors: what proportion of staff had experienced or witnessed verbal or physical violence, whether age and experience sufficed to cope with an aggressive behavior, their perception of severity, associated degree of stress, whether the violence was officially reported, and the level of staff training in dealing with violence. Some of the findings are displayed in the form of tables.

3. Results

From staff members approached, a total of 124 responded to the questionnaire: 62 physicians (50.0%), 44

nurses/midwives (35.48%), and 18 attendants (14.52%). Divided into the category of gender: 79 male (63.7%) and 45 female (36.3%) respondents. Table 1 presents the total numbers and percentage of respondents among different professions, who reported encountering aggressive incidents in their workplace. These findings indicate, that physicians (96.77%) and nurses/midwives (81.82%) have a consistently high level of exposure. A total of 108 (87.1%) out of 124 staff members experienced aggressive behavior.

An approach to compare age groups, respondents were divided into five age intervals: ≤ 20 ($n = 1$), 21–30 ($n = 67$), 31–40 ($n = 43$), 41–50 ($n = 11$), and ≥ 51 ($n = 2$). According to the low numbers of the first and last age interval, respondents were shift to the next/reverse age group. The variations are seen as followed: in the 21–30-age group 68 staff members and 13 in the 41–50-age group. The vast majority of the respondents were represented in the second [$n = 68$ (54.84%)] and third [$n = 43$ (34.67%)] age interval. Whereas the statistical results, regarding age intervals, and between male and female respondents, who reported encountering aggressive incidents, are presented in Table 2.

Based on international literature,^{11,12} three categories of aggressive behavior were presented and participants were required to indicate if they had experienced and/or witnessed single or multiple sources of violence within 5 years, as indicated in Table 3. Unfortunately, the current perceived level of verbal abuse (total of 46.98%) and threatening behavior (total of 33.56%) seems to be frequent in emergency departments. Accordingly, the data were also examined for experience earned in practice on how to deal with threat of violence, abusive behavior, and physical assault. To facilitate comparisons among experience, respondents were divided into four groups. The length of health service career ranged from under 5 years up to more than 15 years: ≤ 5 ($n = 39$), 6–10 ($n = 39$), 11–15 ($n = 21$), and ≥ 15 ($n = 25$). In the first group 89.74%, in the second 94.87%, in the third 80.95%, and in the last group 92.0% of the respondent's encountered aggressive behavior, respectively. Considerably, there were no substantial differences regarding encountering aggressive behavior between each group. The only statistically significant difference that emerged from the research was in relation to the extent of trauma due to aggressive and violent incidents among health care providers, seen in Table 4. Whereas no trau-

Table 1
Respondents among different professions in the emergency departments reporting experiences of aggression over a five-year period ($n = 124$)

	Experience of aggressive behavior (n)	Non-experience of aggressive behavior (n)	Total (n)
<i>Statistical results among profession</i>			
Physician	60 (96.77%)	2 (3.23%)	62 (100.0%)
Nurse/Midwife	36 (81.82%)	8 (18.18%)	44 (100.0%)
Attendant	12 (66.67%)	6 (33.33%)	18 (100.0%)
Total (n)	108 (87.1%)	16 (12.9%)	124 (100.0%)

Table 2
Statistical results regarding age groups, and between male and female respondents, who reported encountering aggressive incidents (male $n = 79$; female $n = 45$)

Age interval	Experience of aggressive behavior (n)			
	Male ($n = 79$)	Total male (n)	Female ($n = 45$)	Total female (n)
<i>Statistical results of valid response</i>				
21–30	35 (81.4%)	43 (100.0%)	20 (80.0%)	25 (100.0%)
31–40	24 (85.72%)	28 (100.0%)	11 (73.33%)	15 (100.0%)
41–50	4 (50.0%)	8 (100.0%)	4 (80.0%)	5 (100.0%)
Total (n)	63 (79.75%)	79 (100.0%)	35 (77.78%)	45 (100.0%)

Table 3

Experiencing and/or witnessing single or multiple sources of aggressive/violent incidents in the emergency departments over a five-year period ($n = 298$)

	Verbal abuse (n)	Threatening behavior (n)	Physical assault (n)	Total (n)
<i>Statistical results of valid response</i>				
Experience of	72 (50.7%)	48 (33.8%)	22 (15.5%)	142 (100.0%)
Witness of	68 (43.59%)	52 (33.33%)	36 (23.08%)	156 (100.0%)
Total (n)	140 (46.98%)	100 (33.56%)	58 (19.46%)	298 (100.0%)
<i>Category of aggressive/violent incidents ($n = 304$)</i>				
<i>Verbal abuse (n)</i>				
Screaming/shouting				88 (28.95%)
Calling of names/swearing				62 (20.39%)
Verbal threats				62 (20.39%)
<i>Threatening behavior (n)</i>				
Rising fists				56 (18.42%)
<i>Physical assault (n)</i>				
Slapping/kicking/grabbing				28 (9.21%)
Use of blunt force/sharp objects				6 (1.98%)
Use of weapon				2 (0.66%)
Total (n)				304 (100.0%)

matic reactions shown in the majority (42.65%) of the respondents 38.24% admitted to psychological problems ever since.

Incidents took place in hospital environments, like examination and treatment rooms (39.28%) and waiting rooms of the emergency departments (36.9%), followed by triage (11.9%), and arenas (10.71%). The respondents indicated that dissatisfaction with waiting/care time (57.2%) has been causally implicated in violent incidents. Furthermore, a related factor was alcohol abuse and illicit drug use (25.7%), and economic problems (17.1%).

Of equal interest is the fact that, in spite of the frequency of violent incidents (87.1%), verbal abuse and threatening behavior was never or rarely reported officially (75.73%). Emphasis on reporting such incidents (22.33%) appeared to involve physical violence. Nevertheless, adequate different types of systems, concerning evidence and reducing the risk of violence, are missing in the majority (28.57%), respectively. During a violent incident, the respondents sought support mainly from un-trained security officers (41.67%), and from attendants (25.0%). It is also a matter of concern, that only 4 (3.23%) respondents in this study indicated, that they had training in the management of aggression and violence. Accordingly, 114 respondents (91.93%) expressed a desire for specialized training to cope with violent incidents.

Table 4

Extent of trauma due to aggressive/violent incidents among health care providers ($n = 136$)

<i>Statistical results of valid response</i>	
Non-traumatic reaction	58 (42.65%)
Psychological trauma	52 (38.24%)
Minor trauma	20 (14.71%)
Life-threatening injuries	6 (4.40%)
Total	136 (100.0%)

Table 5

Short profile of assailants (n)

<i>Demographic statistical results of valid response</i>	
<i>Gender ($n = 137$)</i>	
Male	107 (78.1%)
Female	30 (21.9%)
Total	137 (100.0%)
<i>Assailant ($n = 124$)</i>	
Patient	8 (6.45%)
Relative	80 (64.52%)
Patient and relative	36 (29.03%)
Total	124 (100.0%)
<i>Age (years) ($n = 146$)</i>	
≤20	4 (2.74%)
21–30	76 (52.05%)
31–50	60 (41.1%)
≥50	6 (4.11%)
Total	146 (100.0%)
<i>Education ($n = 186$)</i>	
Analphabet	40 (21.51%)
Primary School	78 (41.94%)
Grammar School	40 (21.51%)
University	28 (15.04%)
Total	186 (100.0%)

There is evidence, that health care providers are likely to encounter aggressive behavior from relatives, than from patients themselves; Table 5 presents a short profile of assailants. Contrary to expectation, the aggressors cannot be identified to a certain kind of type, such as age and education, but can be directed to gender (male = 78.1%).

4. Discussion

Abusive behavior, threat of violence, and physical assault are common in the health care sector, and appear

to be of a greater problem for certain social and healthcare staff than for comparable professions.^{13–15} Nevertheless, there is evidence, that physicians¹⁶ and nurses^{17,18} are more likely to encounter aggressive behavior (Table 1) – in particular by men (Table 5).^{7,19} In general, more than three fourths of the represented (87.1%) staff members in this study encountered aggressive incidents in their workplace (Table 3), respectively. Such high levels of verbal aggression (total of 46.98%) and threatening behavior (33.56%) indicate, that it has become a normalized feature of this type of environment. Anecdotal evidence and other research suggest that less experienced health care providers are more likely to be victims of violence.^{20–22} Considerable, the present study produced no evidence to support this suggestion. Furthermore, there were no substantial differences between age intervals, and between male and female staff members (Table 2). In addition, recognizing the serious immediate and future long-term implications for health, psychological and social development that violence represents for individuals, families, communities, and countries is of importance.¹⁰ The extent of trauma due to violence among staff members in the workplace evaluated in this research can be seen in Table 4.

The primary responsibility is, therefore, to prevent aggressive behavior in health care settings,^{23–25} as the majority of respondents indicated, that dissatisfaction with waiting/care time (57.2%) has been causally implicated in violent incidents, and the location of these occurrences have most been taken place in examination/treatment rooms (39.28%) and waiting rooms (36.9%).

According to Ayrancı et al., “in a developing nation such as Turkey with an aggregate health setting occupancy rate of nearly 100%, such issues as lack of beds, overcrowding, patient backups, not being in control of the situation, not having a national health system, use of the emergency department, or secondary health settings, skimpy funding of primary, secondary, or tertiary health care services, and insufficient staffing, would seem to encourage violence”.¹² Nevertheless, violence in health care settings has its roots not in culture or in different systems of health care.^{26,27} Employers in the health care sector have a duty to provide adequate safety measures and training to reduce risks and ensure, where possible, the health, safety, and welfare of their staff members. Respondents in this survey perceived the response of hospital management and the legal system as inadequate, yet there is growing consensus that violent experiences in the workplace are significantly under-reported (75.73%).^{11,28} Factors influencing the low levels of reporting are: believing management will not support staff, that oneself is to blame, that you will be seen as incompetent, and the belief that verbal abuse and threats are part of the job.²³

But, violence and aggressive behavior are not just a criminal justice problem¹³ or just one involving aberrant behavior attributable to alcohol, drugs, respectively presented with 25.7%, or mental illness, most violent incidents within the hospital setting contain an intentional and,

therefore, aggressive element, but it is often predictable and preventable.^{29–31} Aggressive interactions involve interpersonal-communication and in some incidents staff projecting themselves negatively may consciously or unconsciously inflame and worsen difficult situations. The personal safety is the most important factor; minimizing client provocation will, without doubt, reduce one's risk of being assaulted in the emergency department.²⁸ Accordingly, 114 respondents (91.93%) expressed a desire for specialized training to cope with violent incidents. Even at the current perceived level of aggression most staff members are conscious that they lack many of the communication skills necessary to defuse a potentially violent situation.³² Security and workshops on violence prevention strategies were felt to be the most useful potential interventions.^{33–35}

This was a small study in hospital settings in Bolu, but the response generally echoed evidence from other published literature in Turkey.^{36–38} This current study is limited to the fact, that hospital settings are not offering a *Violence Incident Report Form*^{20,39}, a list designed to simplify the registration of violent events. It contributes to definitional difficulties as well as problems in comparing research findings and using such findings to make work environments safe. Obviously, quantitative research is warranted to determine how widespread or frequent the problem actually is in Turkey. Within this context, the number of articles examining factors related to aggression and violence in hospital settings have increased,^{12,40,41} but so far, it is also evident from this research that it is a serious problem, and it is not something to be lightly dismissed. And fairly clear views about what needs to be done did emerge from this research.

5. Conclusion

The general concept of aggression and violence towards health care providers has become increasingly complex and high profile within the health system. Incidents, which had an impact on the safety and welfare of individuals, tend to be either endured or dealt with directly by the victims, or managed by the powers that be with their own dynastic aims in mind, a situation that is often largely true within the culture of Turkey. However, within modern society steps must be taken to eliminate risk in all its manifestations to access control. Consequently, managers, the organization as a whole, and the governmental authority must take violence, threat, and verbal abuse against staff members seriously, and should consult independent respected professionals in order to make risk assessments. A clear definition of violence, which takes into account severity, perceptions, and professional discretion in reporting and dealing with incidents, and counseling for affected staff members are relevant. The risk management process should also enable the optimum level of care to be given to a client and their relatives, and involves the assessment of risk relating to patient care, care systems, and the environment of care. A balance needs to be struck between

maintaining patients' rights and protecting potential victims of violent behavior. An 'ethical perspective' in drawing up any legislation to deal with this issue would undoubtedly offer a more collaborative and balanced view in this respect. In order to address this theme adequately the necessary resources, such as federal guidelines for a *Violence Incident Report Form*, data collection, analysis, and interpretation, will have to be established and have to be made available to recruit, train, and supervise medical staff and working security devices, as well as to provide enough staff members to cope with increased demand, especially in the emergency departments. Recognition of impending trouble, communication skills to defuse trouble, physical training in situations where reporting has identified that physical violence is a problem, provision of physical skills training with the skills that staff actually need, monitoring the effectiveness of training through on-going reporting, pursuit of litigation against aggressors, working agreement with police and security, protocols in place for the management of acutely disturbed patients to include rapid tranquillization is essential. Finally, attention needs to be paid to public education by using the influence of reporting by the media, and focus public opinion into one of demanding change from the national initiatives and by medical health services management. No doubt, it will be useful and necessary to look at strategies from other parts of the world,^{42–51} which aim to deal with the problem.

Acknowledgement

Authors would like to express our appreciation and gratitude to the staff of the University Hospital, the State Hospital, the Hospital of Gynecology and Pediatrics, and the Health Clinic, of the province of Bolu, Turkey, for their participation in this study.

References

- WHO. Highlights on health in Greece. WHO Regional Office for Europe, Press Office. Draft, 1998. p. 1–41.
- UNDP. Human developments report 1996. Oxford University Press. 1996.
- WHO. Health for all targets: the health policy for Europe. WHO Regional Office for Europe, Press Office. Updated edition, 1993.
- OECD. The reform of health care: a comparative analysis of seventeen OECD countries. Organization for Economic Co-operation and Development, Press Office. Health Policy Studies No.5. 1994.
- Schneider M. Gesundheitssysteme im internationalen Vergleich. BASYS Ausgabe 1994; 1995.
- Ferns T, He D. Terminology, stereotypes and aggressive dynamics in the accident and emergency department. *Acad Emerg Nurs* 2005;13:238–46.
- Zernike W, Sharpe P. Patient aggression in a general hospital setting: do nurses perceive it to be a problem? *Int J Nurs Pract* 1998;4(2):126–33.
- Ryan D. Aggression and violence – a problem in Irish accident and emergency departments? *J Nurs Management* 2006;14:106–15.
- Research Perspectives. Violence against social care staff: qualitative research. National Taskforce, Press Office. Fact Sheet; 2000. p. 1–43.
- Holden RJ. Aggression against nurses. *The Aus Nurs J* 1985;15:44–8.
- Fernandes ChMB, Bouthillette F, Raboud JM, Bullock L, Moore CF, Christenson JM, et al. Violence in the emergency department: a survey of health care workers. *CMAJ* 1999;161(10):1245–8.
- Ayrancı Ü, Yenilmez Ç, Balcı Y, Kaptanoğlu C. Identification of violence in Turkish health care settings. *J Interpers Violence* 2006;21(2):276–96.
- National Task Force. Violence against health care providers. Fact Sheet: National Task Force, Press Office; 2006. p.1–8.
- Winstanley S, Whittington R. Aggression towards health care staff in a UK general hospital: variation among professions and departments. *J Clin Nurs* 2004;13:3–10.
- Whitley GG, Jacobson GA, Gawrys MT. The impact of violence in the health care setting upon nursing education. *J Nurs Educ* 1996;35(5):211–8.
- Wyatt JP, Watt M. Violence towards junior doctors in accident and emergency departments. *J Accid Emerg Med* 1995;12(1):40–2.
- Presley D, Robinson G. Violence in the emergency department: nurses contend with prevention in the health care arena. *Nurs Clin North Am* 2002;37(1):161–9.
- Lynch J, Appelboam R, McQuillan PJ. Survey of abuse and violence by patients and relatives towards intensive care staff. *Anaesthesia* 2003;58(9):893–9.
- Ferns T, Cork A, Rew M. Personal safety in the accident and emergency department. *Brit J Nurs* 2005;14(13):725–30.
- Rose M. A survey of violence toward nursing staff in one large Irish accident and emergency department. *J Emerg Nurs* 1997;23(3):214–9.
- Saines JC. Violence and aggression in A& E: recommendations for action. *Accid Emerg Nurs* 1999;7(1):8–12.
- Lyneham J. Violence in New South Wales emergency departments. *Aust J Adv Nurs* 2000;18(2):8–17.
- DH.A safer place: combating violence against social care staff. Department of Health, Press Office. Fact Sheet; 2006. p. 1–74.
- Vanderslott J. A study of incidents of violence towards staff by patients in an NHS trust hospital. *J Psychiat Mental Health Nursing* 1998;5:291–8.
- Warshaw LJ. Workplace violence: preventive and interventional strategies. *J Occup Environ Med* 1996;38(10):993–1006.
- Morgan S. The problems of aggression and violence for health-care staff. *Prof Nurs* 2001;17(2):107–10.
- Brennan W. We don't have to take this: dealing with violence at work. *Nurs Stand* 2000;14(28 Suppl.):3–17.
- Ferns T. Under-reporting of violent incidents against nursing staff. *Nurs Stand* 2006;20(40):41–5.
- Grenyer BFS, Ilkiw-Lavalle O, Biro Ph, Middleby-Clements J, Comminos A, Coleman M. Safer at work: development and evaluation of an aggression and violence minimization program. *Aust New Zealand J Psychiat* 2004;38:804–10.
- Duxbury J. An evaluation of staff and patient views of and strategies employed to manage inpatient aggression and violence on one mental health unit: a pluralistic design. *Psychiatr Ment Health Nurs* 2002;9(3):325–37.
- Arnetz JE, Arnetz BB. Implementation and evaluation of a practical intervention program for dealing with violence towards health care workers. *J Advanced Nurs* 2000;31(3):668–80.
- Lee F. Violence in A& E: the role of training and self-efficacy. *Nurs Stand* 2001;15(46):33–8.
- Peek-Asa C, Cubbin L, Hubbell K. Violent events and security programs in California emergency departments before and after the 1993 Hospital Security Act. *J Emerg Nurs* 2002;28(5):420–6.
- Rew M, Ferns T. A balanced approach to dealing with violence and aggression at work. *Brit J Nurs* 2005;14(4):227–32.
- Forster JA, Petty MT, Schleiger C, Walters HC. Know workplace violence: developing programs for managing the risk of aggression in the health care setting. *MJA* 2005;183(7):357–61.
- Bakıcı K. İşyerinde cinsel taciz (Sexual harassment in the workplace). PhD Thesis. İstanbul Teknik Üniversitesi. 1998. p. 1–78.
- Güngör S. Sağlık sektöründe cinsel taciz mağdur araştırması (Evaluation of sexual harassment victims in the health care sector). Sosyal

- Bilimler Anabilim Dalı Master Thesis. İ.Ü. Adli Tıp Enstitüsü. 1999. p. 1–85.
38. Durgun B. Hemşirelikte cinsel taciz (Sexual harassment against nurses). Sosyal Bilimler Anabilim Dalı Master Tezi. İ.Ü. Adli Tıp Enstitüsü 1998. p. 1–48.
 39. OSHA. Preventing workplace violence for health care & social service workers. Occupational Safety and Health Administration, US Department of Labor. Guidelines; 2004. p.1–47.
 40. Hancı H, Boz B, Demirkan Ö, Kılıçcıoğlu B, Yağmur F. Acil servislerde şiddete maruz kalan sağlık personeli – anket çalışması (Medical personal subjected to violence in the emergency department – enquiry study). *Acil Tıp Dergisi* 2003;3(2):16–20.
 41. Ayrancı Ü, Yenilmez Ç, Günay Y, Kaptanoğlu C. Çeşitli sağlık kurumlarında ve sağlık meslek gruplarında şiddete uğrama sıklığı (Frequency of being exposed to violence in various health institutions and health profession groups). *Anadolu Psikiyatri Dergisi/Anatolian J Psychiat* 2002;3(3):147–54.
 42. Benveniste KA, Hibbert PD, Runciman WB. Violence in health care: the contribution of the Australian Patient Safety Foundation to incident monitoring and analysis. *MJA* 2005;183(7):348–51.
 43. Jackson D, Clare J, Mannix J. Who would want to be a nurse? Violence in the workplace – a factor in recruitment and retention. *J Nurs Manage* 2002;10:13–20.
 44. Leather R. Workplace violence: scope, definition, and global context. In: Cooper C, Swanson N, editors. *Workplace violence in the health sector state of the art*. Geneva: International Labour Office; 2002. p. 3–18.
 45. Mayhew C, Chappell D. The occupational violence experience of some Australian health workers: an exploratory study. *J Occup Health Safety* 2003;19(6):3–43.
 46. Di Martino V. *Workplace violence in the health sector. Country case studies: Brazil, Bulgaria, Lebanon, Portugal, South Africa, Thailand, and an additional Australian study*. Geneva: International Labour Office; 2002. <http://www.icn.ch/SynthesisReportWorkplaceViolenceHealthSector.pdf>.
 47. NSW Health. *Zero tolerance: response to violence in the NSW Health workplace. Policy and framework guidelines*. NSW Department of Health; 2003. http://www.health.nsw.gov.au/pubs/z/pdf/zero_tol_guide.pdf.
 48. Henry J, Ginn GO. Violence prevention in healthcare organizations within a total quality management framework. *J Nurs Adm* 2002;32:479–86.
 49. Keely BR. Recognition and prevention of hospital violence. *Dimens Crit Care Nurs* 2002;21:236–41.
 50. Walters H, Kay R. Towards a compassionate controls strategy. *Aust Nurs J* 2004;12(4):21–2.
 51. International Labour Office (ILO), International Council of Nurses (ICN), World Health Organization (WHO), Public Services International (PSI). Guidelines on workplace violence in the health sector. Comparison of major known national guidelines and strategies: United Kingdom, Australia, Sweden, USA (OSHA and California). Geneva: Joint Program on Workplace Violence in the Health Sector; 2003. http://www.icn.ch/SewWorkplace/WPV_HS_ComparisonGuidelines.pdf.